

CONSTANT ACCELERATION FORMULAE



$$V = u + at$$

$$S = ut + 0.5at^2$$

$$S = vt - 0.5at^2$$

$$S = \frac{(u + v) \times t}{2}$$

$$v^2 = u^2 + 2as$$

v = final velocity (m/s)
 u = initial velocity (m/s)
 a = acceleration (m/s²)
 s = displacement (m)
 t = time (s)

TORQUE

$$T = F \times d \times \sin(\theta)$$



T = Torque (Nm)
 F = Force applied on lever (N)
 d = Distance that the force is applied from the axis of rotation (m)
 θ = Angle between force vector and distance vector, sin(θ) is equal to 1 if perpendicular

NEWTON'S 2ND LAW OF MOTION



$$\Sigma F = m \times a$$

ΣF = net force acting on an object (N)
 m = mass of object (kg)
 a = acceleration of object (m/s²)


MECHANICAL ENERGY

$$ME = U_s + U_g + KE$$



ME = mechanical energy (J)
 U_s = spring potential energy, a.k.a elastic potential energy (J)
 U_g = gravitational potential energy (J)
 KE = kinetic energy (J)

MOMENTUM



$$p = mv$$

p = Momentum (kg x m/s)
 m = Mass (kg)
 v = Velocity (m/s)

GRAVITATIONAL POTENTIAL ENERGY

$$U_g = mgh$$



U_g = gravitational potential energy (J)
 m = Mass of object (kg)
 g = Acceleration due to gravity = 9.8 m/s² = gravitational field strength = 9.8 N/kg
 h = Vertical height above a reference level (m)

TEMPERATURE CHANGE



$$Q = m \times C \times (T_{\text{final}} - T_{\text{initial}})$$

Q = heat energy applied (J)
m = mass of object (kg)
c = specific heat capacity of object (J/kg/K)
T = initial and final temperatures of object (K)

FORCE ON A MOVING CHARGE



$$F = qvb$$

F = Magnetic Force
q = Charge of the particle
v = Velocity of the charged particle
b = Magnetic Field Strength


OHM'S LAW



$$V = IR$$

V = Voltage (volts)
I = Current (amps)
R = Resistance (ohms)

ELECTRICAL POWER



$$P = IV = I^2R = \frac{V^2}{R}$$

P = Power
V = Voltage (volts)
I = Current (amps)
R = Resistance (ohms)

KINETIC ENERGY



$$KE = \frac{1}{2} mv^2$$

KE = Kinetic energy of object (J)
m = Mass of object (kg)
v = Velocity of object (m/s)

GRAVITATIONAL FORCE



$$F = \frac{G \times m_1 \times m_2}{r^2}$$

F = gravitational force of attraction from the center of mass of the two objects (N)
G = gravitational constant 6.674×10^{-11} Nm kg
m1 = mass of one of the objects (kg)
m2 = mass of the other object (kg)
r = distance between the center of the mass of the two objects (m)

PRESSURE



$$P = \frac{F}{A}$$

P = Pressure (N/m or Pascals)
F = Force applied on a surface (N)
A = Area of surface (m²)

ARCHIMEDES PRINCIPLE



$$F_b = F_{g \text{ of object}} = F_{g \text{ of fluid displaced}} = \rho_{\text{fluid}} \times V_{g \text{ of fluid displaced}} \times g$$

F_b = Force of buoyancy (N)
F_{g of object} = Weight of object (N)
F_{g of fluid displaced} = Weight of fluid displaced (N)
ρ_{fluid} = Density of fluid (kg/m³)
V_{fluid displaced} = Volume of fluid displaced by the object in the fluid (m³)
g = gravitational field strength, a.k.a acceleration due to gravity = 9.8 N/kg = 9.8 m/s²

Don't Miss out on our Free Medical School Admissions Webinar!

Cover the Admissions Milestones with Current Med Students

Confused by the Medical Schools Admission Process? Want tips from a Current Medical Student on the Admissions Milestones and Criteria?

Here at GradReady we're running a free Medical School Admissions Webinar which will provide students with a breakdown of the changes and key milestones.



This webinar will cover the admission process of medical schools in 5 states: VIC, NSW, QLD, WA & SA.

We'll also cover portfolios in all these webinars with advice for UNDS, UNDF, UOW & MQ.


This webinar will highlight the admissions milestones and the GEMSAS admissions process (including details on preferencing) and canvass the various admissions criteria for graduate medical schools across Australia. You'll have the opportunity to ask all your burning questions about the long year ahead.

For further details visit:

gradready.com.au/gamsat/events

You can also visit our [FB Page](#) to find booking information and event details.

ELECTROSTATIC FORCE



$$F = \frac{k \times q_1 \times q_2}{r^2}$$

F = electrostatic force acting between two charged objects (N)

k = Coulomb's constant = $9.0 \times 10 \text{ NmC}$

q1 = magnitude of one of the charged object (C)

q2 = magnitude of the other charged object (C)

r = distance between the center of the two charges (m)

WORK



$$W = F \times d \times \cos(\theta)$$

W = Work done (J)

F = Force applied on object (N)

d = Distance object moves due to the application of the force (m)

θ = Angle between force vector and displacement vector, $\cos(\theta)$ is equal to 1 when the force vector is in the same direction as the movement of the object

or

$$W = F \times d$$

W = Work done (J)

F = Component of the force that acts parallel to the distance only (N)

d = Distance object moves due to the application of the force (m)



Average Improvement of **20+** Percentile Points
Over the last **10+** Years

10,000+
Happy Students

10+
Years Experience

Most Advanced
Technology